ADOFTION Information, resources, and access in Northern Ontario





First day of last menstrual period (LMP):

Positive urine pregnancy test date:





Ultrasound may be necessary if:

- · LMP is uncertain
- · LMP was more than 70 days ago
- Ectopic pregnancy is suspected
- Risk factors for ectopic pregnancy

Medication abortion





Day 1: Take anti-progesterone tablet (Mifepristone 200 mg) orally

Day 2: Take prostaglandin tablet (Misoprostol 800 mcg) buccally (2 tablets in each cheek pouch for 30 minutes, then swallow)

Expectations: Painful uterine cramping, heavy vaginal bleeding, passage of tissue +/- nausea, diarrhea, headache

 Vaginal bleeding is heaviest on day 1 and 2, and decreases over 2 weeks **Seek medical attention** if you develop a fever ≥38 °C for more than 3 hours, soak through 1 pad/hour after initial 6 hours, or experience uncontrolled pain

Procedural abortion

Dilation: Provider will insert a "dilator" wand into your cervix to gently stretch open your cervical opening.



Curettage: provider will insert a "curette" tube into your cervical opening and gently suction to empty the contents of your uterus.

Which method is best for me?



| | Medication abortion | Procedural abortion |
|-----------------------|---------------------|---------------------------|
| Location: | Home | Hospital |
| Consciousness: | Aware | Under anesthesia/sedation |
| Duration of bleeding: | Heavier & longer | Lighter & shorter |
| Effectiveness: | 98% | >99% |
| | | |

Support



Action Canada (Planned Parenthood Canada)

1.888.642.2725 ActionCanadaSHR.org



AbortionSupportCollective.ca

PregnancyOptions.info

Accessing abortion





Virtual medical abortion clinic: CircleOfCareHealthConnections.com

Locate a provider: ChoiceConnect.ca



Visit a physician or nurse practitioner at any medical clinic for (a) prescription for a medication abortion, or (b) referral for procedural abortion

Call Options Clinic at Health Sciences North 1.705.671.5265



Adapted from:

Canadian Protocol for the Provision of Medical Abortion, 2020.

Induced Abortion: Updated Guidance During Pandemics and Periods of Social Disruption, 2020.

01/2023

phsd.ca



■ PublicHealthSD

