Vaccine Order Form

phsd.ca tel: 705.522.9200, ext. 490 toll-free: 1.866.522.9200



Orders placed Monday to Friday by noon will be available for pick up the following Monday between 8:30 a.m. and noon or 1 and 4:30 p.m. For all partners outside of Greater Sudbury, your local Public Health office will contact you when your order is available for pick up.

Office information	·	Email orders	
Date	# of HCPs	vaccineorder@phsd.ca	
Office name and phone #		Fax orders	
Office contact and email		1.705.677.9616	

A hard-sided, pre-conditioned cooler, two cool blankets, three ice packs and one temperature monitoring device are required to receive and transport vaccines. Always maintain vaccine temperatures between 2°C and 8°C.							
ROUTINE vaccine	Description	Doses on hand	Doses required				
Adacel [®] /Boostrix [®]	Tetanus, Diphtheria and Pertussis vaccine						
Adacel-Polio [®] /Boostrix-Polio [®]	Tetanus, Diphtheria, Pertussis and Polio vaccine						
Menjugate [®] /NeisVac-C [®]	Meningococcal Conjugate C liquid vaccine						
MMRII [®] /Priorix [®]	Measles, Mumps and Rubella vaccine						
Proquad [®] /Priorix-Tetra [®]	Measles, Mumps, Rubella and Varicella vaccine						
Pediacel [®]	Pertussis, Diphtheria, Tetanus, Polio and Haemophilus influenzae type B vaccine						
Pneumovax 23 [®]	Pneumococcal Polysaccharide vaccine						
Polio [®]	Polio vaccine	1 dose/box					
Prevnar 13 [®]	Pneumococcal Conjugate vaccine - 13 valent	10 doses/box					
Rotarix [®]	Rotavirus vaccine (Rot-1)	10 doses/box					
Td Adsorbed [®]	Tetanus and Diphtheria vaccine 5 doses/box						
Tubersol [®]	Tuberculin Skin testing solution 10 doses/vial						
Varilrix [®] /Varivax III [®]	Varicella vaccine	10 doses per box					
Shingrix [®]	Varicella-Zoster shingles vaccine (aged 65-70 years)	1 /box or 10 /box					
RESPIRATORY Vaccines	Description	0 0	Doses on hand	Doses required			
Fluzone® or Flulaval®	Quadrivalent Influenza; multi-dose vial or prefilled syringe. Age 6 months and older.						
High-Dose Fluzone®	High-Dose Quadrivalent Influenza; prefilled 5 doses/box syringe. Age 65 years and older						
Fluad®	Adjuvanted Trivalent Influenza; prefilled syringe. Age 65 years and older						
Pfizer-BioNTech Comirnaty®	Omicron XBB.1.5 subvariant COVID-19 vaccine. Age 12 years and older	6 doses/vial					
	Omicron XBB.1.5 subvariant COVID-19 vaccine. Age 5 to 12 years of age	10 doses/vial					
	Omicron XBB.1.5 subvariant COVID-19 vaccine. Age 6 months to 5 years of age	10 doses/vial					
Moderna Spikevax® *If available	Omicron XBB.1.5 subvariant COVID-19 vaccine. Age 6 months and older	5 doses/vial					
Novavax® or ancestral Bivalent COVID Arexvy®	*Upon request, if available Respiratory syncytial virus vaccine; LTCH/Elder Care Lodge resident AND age 60 years and older						

HIGH-RISK and SCHOOL vac	cines		
Vaccine name	DOB (YYYY/M/D)	Doses required	Eligibility Criteria
Human Papillomavirus (Gardasil-9 [®])			☐ Grade 7-12 students☐ Men who have sex men ≥ 9 years to < 27 years
Hepatitis A (Avaxim [®] /Havrix [®] /Vaqta®			≥ 1 year with: (please check all that apply) ☐ Chronic liver disease (including Hepatitis B and C) ☐ Persons engaging in intravenous drug use ☐ Men who have sex with men
Hepatitis B (Recombivax HB®/Engerix-B®)			Grade 7-12 students ≥ 0 years with: (please check all that apply) Children <7 years old whose families have immigrated from countries of high prevalence for HBV, and who may be exposed to HBV carriers through their extended families (3 doses) Household and sexual contacts of chronic carriers and acute cases History of a sexually transmitted disease (3 doses) Infants born to HBV-positive carrier mothers: premature infants weighing <2,000 grams at birth (4 doses) premature infants weighing ≥2,000 grams at birth and full/post term infants (3 doses) Intravenous drug use (3 doses) Liver disease (chronic), including hepatitis C (3 doses) Men who have sex with men (3 doses) Multiple sex partners (3 doses) Needle stick injuries in a non-health care setting (3 doses) On renal dialysis or those with diseases requiring frequent receipt of blood products (e.g., haemophilia) (2nd and 3rd doses only)
Haemophilus influenzae type b (Act-HIB [®] /Hiberix [®])			 ☐ Awaiting liver transplants (2nd and 3rd doses only) ≥ 5 years with: (please check all that apply) ☐ Asplenia (functional or anatomic) (1 dose) ☐ Bone marrow or solid organ transplant recipients (1 dose) ☐ Cochlear implant recipients (pre/post implant) (1 dose) ☐ Hematopoietic stem cell transplant (HSCT) recipients (3 doses) ☐ Immunocompromised individuals related to disease/therapy (1 dose) ☐ Lung transplant recipients (1 dose) ☐ Primary antibody deficiencies (1 dose)
Meningococcal B (Bexsero [®])			Age 2 months to 17 years with: (please check all that apply) ☐ Acquired complement deficiencies (e.g., receiving eculizumab) ☐ Asplenia (functional or anatomic) ☐ Cochlear implant recipients (pre/post implant) ☐ Complement, properdin, factor D or primary antibody deficiencies ☐ HIV
Meningococcal C-ACYW135 (Menactra®/Nimenrix®)			 ☐ Grade 7-12 students ☐ Born in or after 1997 Age 9 months to 55 years with: (please check all that apply) ☐ Functional or anatomic asplenia ☐ Complement, properdin, factor D or primary antibody deficiency ☐ Cochlear implant recipient (pre/post implant) ☐ Acquired complement deficiency ☐ HIV
FOR OFFICE USE ONLY: ☐ Are completed, cold chain of a confirmed agreement in plate of the confirmed to partner store of the confirmed to partner store of the confirmed agreement in plate of the confirmed agre	ace for COVI	D-19? x?	e logs of the past two weeks included with the order?